




15019AST Supelco

Astec® CHIROBIOTIC® V2 Chiral HPLC Column

5 µm particle size, L × I.D. 15 cm × 2.1 mm

eCl@ss 32110501



FDS

Similar Products

Conditionnement - SKU	Disponibilité	Prix (EUR)	Quantité
15019AST	Seulement 1 en stock (d'autres en cours d'arrivage) - A PARTIR DE	2,092.00	0

Commandes Bulk?

AJOUTER AU PANIER

Propriétés

Related Categories	Analytical/Chromatography, Astec CHIROBIOTIC Chiral HPLC Columns (Macrocyclic Glycopeptides), Astec CHIROBIOTIC V2 Chiral HPLC Columns, CHIROBIOTIC V2, Chiral HPLC Columns, HPLC Less...
description	HPLC column
packaging	pkg of 1 ea
application(s)	HPLC: suitable
L × I.D.	15 cm × 2.1 mm
matrix	High-purity silica gel particle platform
matrix active group	vancomycin phase
particle size	5 µm
pore size	200 Å
operating pH range	3.5 - 7.0
mode of chromatography	chiral

montrer moins de résultats

Description

General description

Neutral molecules, amides, acids, esters and amines show considerable enantioselectivity on these vancomycin-based CSPs. A wide variety of secondary and tertiary amines have been separated on the CHIROBIOTIC® V in the polar ionic mode. The CHIROBIOTIC® V has demonstrated many of the separation characteristics of protein-based stationary phases, but with exceptional stability and much higher sample capacity. Some chiral analytes have been resolved that have not been reported separated on any other chiral stationary phase. CHIROBIOTIC® V and V2 differ in their bonding chemistry the pore size of the support particle, giving them different selectivity and preparative capacity.

- Bonded phase: Vancomycin
- Operating pH range: 3.5 - 7.0
- Particle diameter: 5, 10 or 16 µm
- Pore size: 100 Å (CHIROBIOTIC® V) or 200 Å (CHIROBIOTIC® V2)

[CHIROBIOTIC FAQs](#)
[CHIROBIOTIC Reference Bibliography](#)
[Chiral Product Literature](#)

Recommended products

Discover LiChropur reagents ideal for **HPLC** or **LC-MS** analysis

Legal Information

Astec is a registered trademark of Sigma-Aldrich Co. LLC

CHIROBIOTIC is a registered trademark of Sigma-Aldrich Co. LLC

Aucune information relative à la sécurité de ce produit est actuellement disponible.

Certificat d'Analyse

8854

[Devis/ Commande de produits Bulk](#)

[FDS](#)

Customers Also Viewed

15021AST Astec® CHIROBIOTIC® V2 Chiral HPLC Column 5 µm particle size, L × I.D. 5 cm × 4.6 mm	14019AST Astec® CHIROBIOTIC® TAG Chiral HPLC Column 5 µm particle size, L × I.D. 15 cm × 2.1 mm	11019AST Astec® CHIROBIOTIC® V Chiral HPLC Column 5 µm particle size, L × I.D. 15 cm × 2.1 mm	15020AST Astec® CHIROBIOTIC® V2 Chiral HPLC Column 5 µm particle size, L × I.D. 25 cm × 2.1 mm	15023AST Astec® CHIROBIOTIC® V2 Chiral HPLC Column 5 µm particle size, L × I.D. 15 cm × 4.6 mm
--	--	--	---	---

Recently Viewed

15018AST Astec® CHIROBIOTIC® V2 Chiral HPLC Column 5 µm particle size, L × I.D. 10 cm × 2.1 mm	14100AST Astec® CHIROBIOTIC® TAG Chiral HPLC Guard Cartridge, 5 µm particle size, L × I.D. 2 cm × 4 mm	14044AST Astec® CHIROBIOTIC® TAG Chiral HPLC Column 5 µm particle size, L × I.D. 25 cm × 21.2 mm	14034AST Astec® CHIROBIOTIC® TAG Chiral HPLC Column 5 µm particle size, L × I.D. 25 cm × 10 mm
---	---	---	---

Protocoles et articles

Articles

Fast LC-MS on Astec CHIROBIOTIC CSPs

Astec CHIROBIOTIC® chiral stationary phases (CSPs) offer several different types of molecular interactions, and separate enantiomers of many analyte classes. Of particular interest are ionic interactions. Craig Aurand
Reporter US Volume 27.2
Keywords: High performance liquid chromatography, Liquid chromatography mass spectrometry, Mass spectrometry, Phase transitions, Separation

Related Content

Supelco HPLC Calculator for Isocratic or Gradient Method Transfer

Calculates conditions for transfer of an isocratic or gradient method from one HPLC column to another. Allows method scaling from microbore through preparative column range. Calculates time and solve...
Keywords: High performance liquid chromatography

Documentation référencée

Did you use this product in your Paper? If so click here.

Enantiomeric impurities in chiral catalysts, auxiliaries, and synthons used in enantioselective syntheses. Part 4 Qui, Haiziao, et al. Tetrahedron Asymmetry 24 (18), 1134-1141, (2013)

[read abstract](#)

High-performance liquid chromatographic enantioseparation of unusual isoxazoline-fused 2-aminocyclopentanecarboxylic acids on macrocyclic glycopeptide-based chiral stationary phases.

László Sipos et. al
Journal of chromatography. A, 1232, undefined (2011-12-20)
The enantiomers of four unusual isoxazoline-fused 2-aminocyclopentanecarboxylic acids were directly separated on chiral stationary phases containing macrocyclic glycopeptide antibiotics teicoplanin (Astec Chirobiotic T and T2), teicoplanin aglycone (...[Read More](#))

[read abstract](#)

New high-performance liquid chromatography method for the determination of (R)-warfarin and (S)-warfarin using chiral separation on a glycopeptide-based stationary phase.

Jana Malakova et. al
Journal of chromatography. B, Analytical technologies in the biomedical and life sciences, 877(27), undefined (2009-8-1)
Warfarin is a well-known anticoagulant agent that occurs in two enantiomers, (R)-(+)-warfarin and (S)-(-)-warfarin. A new liquid chromatography method for the determination of both enantiomers was developed, validated and applied in in vitro studies ...[Read More](#)
Chiral liquid chromatography/tandem mass spectrometry assay to determine that dexpropimexole is not converted to pramipexole in vivo after administered in humans Wei, Dong, et al. Journal of Chromatography. B, Biomedical Sciences and Applications 971, 133-140, (2014)
New chemo-enzymatic approaches for the synthesis of (R)- and (S)-bupropionolol Nagy, Botond, et al. Tetrahedron Asymmetry 25 (18-19), 1316-1322, (2014)

[read abstract](#)



Development and validation of an UFLC-MS/MS method for enantioselectivity determination of d,l-threo-methylphenidate, d,l-threo-ethylphenidate and d,l-threo-ritalinic acid in rat plasma and its application to pharmacokinetic study.

Chenghao Zhang et. al

Journal of chromatography. B, Analytical technologies in the biomedical and life sciences, 1011, undefined (2016-1-12)

A chiral UFLC-MS/MS method was established and validated for quantifying d-threo-methylphenidate (d-threo-MPH), l-threo-methylphenidate (l-threo-MPH), d-threo-ethylphenidate (d-threo-EPH), l-threo-ethylphenidate (l-threo-EPH) and d,l-threo-ritalinic ...[Read More](#)

Enantioselective and sensitive determination of carvedilol in human plasma using chiral stationary-phase column and reverse-phase liquid chromatography with tandem mass spectrometry Jiang, Juanjuan, et al. Journal of Chromatography. B, Biomedical Sciences and Applications 960, 92-97, (2014)

Validation of a chiral liquid chromatography/tandem mass spectrometry method for the determination of pantoprazole in dog plasma Chen, Meixia, et al. Journal of Chromatography. B, Biomedical Sciences and Applications 906, 85-90, (2012)

[read abstract](#)

Development, optimization and validation of a sub-minute analytical enantioselective high performance liquid chromatographic separation for a folic acid precursor in normal phase mode.

Doris Frühauf and Markus Juza

Journal of chromatography. A, 1269, undefined (2012-10-30)

A sub-minute enantioselective normal phase high performance liquid chromatographic (HPLC) method for the analysis of a chiral precursor molecule employed frequently in folic acid syntheses was developed, optimized and successfully validated according...[Read More](#)

Enantiospecific determination of dl-methylphenidate and dl-ethylphenidate in plasma by liquid chromatography/tandem mass spectrometry: Application to human ethanol interactions Zhu, Hao-Jie, et al.

Journal of Chromatography. B, Biomedical Sciences and Applications 879 (11-12), 783-788, (2011)

Chiral derivatives of Butenafine and Terbinafine: synthesis and antifungal activity Fuglseth, Erik, et al. Tetrahedron 65 (47), 9807-9813, (2009)

Enantioselective determination of doxazosin in human plasma by liquid chromatography/tandem mass spectrometry using ovomucoid chiral stationary phase Liu, Ke, et al. Journal of Chromatography. B, Biomedical Sciences and Applications 878 (26), 2415-2420, (2010)

Enantioenriched 1-aryl-2-fluoroethylamines. Efficient lipase-catalysed resolution and limitations to the Mitsunobu inversion protocol Thvedt, Thor H?kon Krane, et al. Tetrahedron 66 (34), 6733-6743, (2010)

Quantitative determination of ondansetron in human plasma by enantioselective liquid chromatography-tandem mass spectrometry Liu, Ke, et al. Journal of Chromatography. B, Biomedical Sciences and Applications 864 (1-2), 129-136, (2008)

Direct and indirect separations of five isomers of Brivanib Alaninate using chiral high-performance liquid chromatography He, Brian Lingfeng, et al. Journal of Chromatography. B, Biomedical Sciences and Applications 875 (1), 122-135, (2008)

[Show more references \(6 remain\)](#)


Produits associés



**Brighter
LC-MS**

LC-MS Grade Solvents and Reagents
for accurate, brilliant results

[LEARN MORE](#)



**Milli-Q® water
purification
solutions**

Ultrapure & Pure Systems
for Every Lab

[Learn More](#)

© 2018 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved. Reproduction of any materials from the site is strictly forbidden without permission. Sigma-Aldrich Products are sold exclusively through Sigma-Aldrich, Inc. [Site Use](#)
[Terms](#) | [Privacy](#)