



20019AST Supelco

Astec® CYCLOBOND® I 2000 Chiral HPLC Column

5 µm particle size, L × I.D. 15 cm × 2.1 mm

eCl@ss 32110501

FDS

Similar Products

Conditionnement - SKU	Disponibilité	Prix (EUR)	Quantité
20019AST	Seulement 1 en stock (d'autres en cours d'arrivage) - A PARTIR DE	1,626.00	0

Commandes Bulk?

AJOUTER AU PANIER

Propriétés

Related Categories	Analytical/Chromatography, Astec CYCLOBOND Chiral HPLC Columns (Native and Derivatized Cyclodextrins), CYCLOBOND I 2000, Chiral HPLC Columns, HPLC Less...
packaging	pkg of 1 ea
application(s)	HPLC: suitable
L × I.D.	15 cm × 2.1 mm
matrix	silica particle platform
matrix active group	cyclodextrin, beta- phase
particle size	5 µm
pore size	100 Å
mode of chromatography	chiral

[montrer moins de résultats](#)

Description

General description
CYCLOBOND I 2000 is bonded with β-cyclodextrin by a patented process to produce a stable matrix with the cyclodextrin arranged in such a way as to retain its most valuable property of forming inclusion complexes. This allows them to effect numerous chemical separations by selectively including into their cavities a wide variety of organic molecules. Non-inclusion type separations are also possible with the polar organic mode for a wide variety of molecule types.

- Bonded phase: Underivatized, native β-cyclodextrin

Resources:
[CYCLOBOND FAQs](#)
[CYCLOBOND Reference Bibliography](#)
[Chiral Product Literature](#)

Recommended products
Discover LiChropur reagents ideal for **HPLC** or **LC-MS** analysis

Legal Information
Astec is a registered trademark of Sigma-Aldrich Co. LLC
CYCLOBOND is a registered trademark of Sigma-Aldrich Co. LLC

Informations Sécurité

Aucune information relative à la sécurité de ce produit est actuellement disponible.

Documents

Certificat d'Analyse

6341

Devis/ Commande de produits Bulk

FDS

Customers Also Viewed

 20024AST Astec® CYCLOBOND® I 2000 Chiral HPLC Column 5 µm particle size, L × I.D. 25 cm × 4.6 mm	 20018AST Astec® CYCLOBOND® I 2000 Chiral HPLC Column 5 µm particle size, L × I.D. 10 cm × 2.1 mm	 20013-U Carbotrap® 217 stainless steel TD tube, O.D. × L 1/4 in. × 3 1/2 in., preconditioned, pkg of 10 ea	 20012-U Carbosieve® SIII stainless steel TD tube, O.D. × L 1/4 in. × 3 1/2 in., preconditioned, pkg of 10 ea	 20011-U Tenax® GR stainless steel TD tube, O.D. × L 1/4 in. × 3 1/2 in., preconditioned, pkg of 10 ea
---	---	---	---	--

Recently Viewed

 20018AST Astec® CYCLOBOND® I 2000 Chiral HPLC Column 5 µm particle size, L × I.D. 10 cm × 2.1 mm	 20013-U Carbotrap® 217 stainless steel TD tube, O.D. × L 1/4 in. × 3 1/2 in., preconditioned, pkg of 10 ea	 20012-U Carbosieve® SIII stainless steel TD tube, O.D. × L 1/4 in. × 3 1/2 in., preconditioned, pkg of 10 ea	 20011-U Tenax® GR stainless steel TD tube, O.D. × L 1/4 in. × 3 1/2 in., preconditioned, pkg of 10 ea	 20010-U Tenax® TA stainless steel TD tube, O.D. × L 1/4 in. × 3 1/2 in., preconditioned, pkg of 10 ea
---	---	---	--	--

Protocoles et articles

Articles

[Analysis of Isoprenoid Pathway Metabolites by LC-MS | Biofiles, Vol. 8, No. 17](#)

Isoprenoid pathway metabolites such as isoprenoid phosphates and pyrophosphates are central biological intermediates leading to sterols, dolichols, ubiquinones, prenylated natural products and proteins. ...
Rudolf Köhling, Roland Meier, Bernhard Schönenberger and Roland Wohlgenuth

Sigma-Aldrich®, Buchs, Switzerland

Biofiles, Vol. 8, No. 17

Keywords: Buffers, Chromatography, High performance liquid chromatography, Individual protein Immunoprecipitation, Mass spectrometry, Metabolites, Separation, Solvents

Related Content

[Supelco HPLC Calculator for Isocratic or Gradient Method Transfer](#)

Calculates conditions for transfer of an isocratic or gradient method from one HPLC column to another. Allows method scaling from microbore through preparative column range. Calculates time and solve...

Keywords: High performance liquid chromatography

Documentation référencée

[Did you use this product in your Paper? If so click here.](#)

High-performance liquid chromatography of diastereomeric flavanone glycosides in Citrus on a β -cyclodextrin-bonded stationary phase (Cyclobond I) Krause, Martin; Galensa, Rudolf Journal of Chromatography A 588 (1-2), 41-45, (1991)

[read abstract](#)

Analytical method development for directed enzyme evolution research: a high throughput high-performance liquid chromatography method for analysis of ribose and ribitol and a capillary electrophoresis method for the separation of ribose enantiomers.

Baoguo Sun et. al

Journal of chromatography. A, 1271(1), undefined (2012-12-15)

Analytical methods were developed for a directed enzyme evolution research programme, which pursued high performance enzymes to produce high quality L-ribose using large scale biocatalytic reaction. A high throughput HPLC method with evaporative light scattering detection...[Read More](#)

[read abstract](#)

Separation and characterization of underivatized oligosaccharides using liquid chromatography and liquid chromatography-electrospray ionization mass spectrometry.

Ying Liu et. al

Journal of chromatography. A, 1079(1-2), undefined (2005-7-26)

Native cyclodextrin-based columns are particularly useful for the analysis of oligosaccharides because the retention of these carbohydrates is based mainly on the hydrogen bonding interactions of oligosaccharide hydroxyl groups with the stationary phase...[Read More](#)

[read abstract](#)

The enantioselective determination of chlorpheniramine and its major metabolites in human plasma using chiral chromatography on a beta-cyclodextrin chiral stationary phase and mass spectrometric detection.

Karen M Fried et. al

Journal of pharmaceutical and biomedical analysis, 27(3-4), undefined (2002-1-5)

A sensitive enantioselective high-performance liquid chromatographic (HPLC) method has been developed for the simultaneous determination of plasma concentrations of (-)(R)- and (+)(S)-chlorpheniramine (CP) and their metabolites, desmethyl-chlorpheniramine...[Read More](#)

[read abstract](#)

High-performance liquid chromatographic separation and on-line mass spectrometric detection of saturated and unsaturated oligogalacturonic acids.

Thomas Stoll et. al

Carbohydrate research, 337(24), undefined (2002-12-21)

A method for the simultaneous determination of saturated and unsaturated oligogalacturonic acids up to degree of polymerization (dp) of 7 by high-performance liquid chromatography (HPLC) is presented. For this purpose, a Cyclobond I 2000 column and a...[Read More](#)



[read abstract](#)

Separation of chiral furan derivatives by liquid chromatography using cyclodextrin-based chiral stationary phases.

Xinxin Han et. al

Journal of chromatography. A, 1063(1-2), undefined (2005-2-11)

The enantiomeric separation of a set of 30 new chiral furan derivatives has been achieved on native and derivatized beta-cyclodextrin stationary phases using high performance liquid chromatography (HPLC). The hydroxypropyl-beta-cyclodextrin (Cyclobon...[Read More](#)

[read abstract](#)

Separation of stereoisomers of sertraline and its related enantiomeric impurities on a dimethylated beta-cyclodextrin stationary phase by HPLC.

R Nageswara Rao et. al

Journal of pharmaceutical and biomedical analysis, 50(3), undefined (2009-6-9)

A reversed-phase high-performance liquid chromatographic (HPLC) method was developed and validated for evaluating the chiral discrimination ability of CYCLOBOND I 2000 DM chiral stationary phase (CSP) towards sertraline and its related enantiomers. T...[Read More](#)

[read abstract](#)

Screening approach, optimisation and scale-up for chiral liquid chromatography of cathinones.

R Wimal H Perera et. al

Journal of chromatography. A, 1269, undefined (2012-11-24)

Screening approaches adopted in pharmaceutical companies for chiral LC method development may be quite complicated and sophisticated in order to guarantee a high success rate. However in other environments it may be of more value to assess how simple...[Read More](#)

[read abstract](#)

Assessment of chiral stationary phases for suitability for combined enantiomeric impurity/related substances assays.

R Wimal H Perera and W John Lough

Journal of chromatography. A, 1218(48), undefined (2011-11-2)

Chiral stationary phases (CSP) for LC had a major impact on pharmaceutical R&D when they first became commercially available in the 1980s. Even although the use of CSP in pharmaceutical R&D is now very much a mature area, there is still scope for usi...[Read More](#)

[read abstract](#)

Separation of oxazepam, lorazepam, and temazepam enantiomers by HPLC on a derivatized cyclodextrin-bonded phase: application to the determination of oxazepam in plasma.

Chuong Pham-Huy et. al

Journal of biochemical and biophysical methods, 54(1-3), undefined (2003-1-25)

The enantioselective high-performance liquid chromatography (HPLC) of three racemic 3-hydroxybenzodiazepines, oxazepam (Oxa), lorazepam (Lor), and temazepam (Tem), is a difficult operation because of the spontaneous chiral inversion in polar solvent....[Read More](#)

[read abstract](#)

HPLC and 1H-NMR study of chiral recognition in some thromboxane antagonists induced by beta-cyclodextrin.

A F Casy et. al

Journal of pharmaceutical and biomedical analysis, 9(10-12), undefined (1991-1-1)

The interaction of beta-cyclodextrin with a series of structurally related chiral thromboxane antagonists was investigated using NMR and RP-HPLC. HPLC studies used both a cyclodextrin bonded phase (Cyclobond I), and beta-cyclodextrin as a mobile phas...[Read More](#)

Liquid chromatographic resolution of enantiomers of deltahedral carborane and metallaborane derivatives Plesek, Jaromir, et al. Journal of Chromatography A 626 (2), 197-206, (1992)

[read abstract](#)

Chiral stationary phases in HPLC for the stereoselective determination of methadone.

S Rudaz and J L Veuthey

Chirality, 11(4), undefined (1999-5-29)

An extensive study of the behavior of three chiral stationary phases (CSP) is presented for the stereoselective determination of methadone. The following chromatographic columns were selected: a cellulose, Chiralcel OJ; a modified cyclodextrin. Cyclo...[Read More](#)

[read abstract](#)

High-performance liquid chromatographic enantioseparation of bicyclic 1,3-amino alcohols.

A Péter et. al

Journal of chromatography. A, 919(1), undefined (2001-7-19)

Different high-performance liquid chromatographic methods were developed for the separation and identification of enantiomers of diendo- and diexo-3-aminobicyclo[2.2.1]heptane-2-methanol and diendo- and diexo-3-amino-bicyclo[2.2.1]hept-5-ene-2-methan...[Read More](#)

Chiral separations of ?-blocking drug substances using chiral stationary phases Ekelund, Jens, et al. Journal of Chromatography A 708 (2), 253-261, (1995)


[Show more references \(20 remain\)](#)

Produits associés



Brighter LC-MS
LC-MS Grade Solvents and Reagents for accurate, brilliant results

[LEARN MORE](#)



Milli-Q® water purification solutions
Ultrapure & Pure Systems for Every Lab

[Learn More](#)

© 2018 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved. Reproduction of any materials from the site is strictly forbidden without permission. Sigma-Aldrich Products are sold exclusively through Sigma-Aldrich, Inc. [Site Use Terms](#) | [Privacy](#)