



20722AST Supelco

Astec® CYCLOBOND® I 2000 DMP Chiral HPLC Column

5 µm particle size, L × I.D. 10 cm × 4.6 mm

eCl@ss 32110501

FDS

Similar Products

Conditionnement - SKU	Disponibilité	Prix (EUR)	Quantité
20722AST	Seulement 2 en stock (d'autres en cours d'arrivage) - A PARTIR DE	1,636.00	0

Commandes Bulk?

AJOUTER AU PANIER

Propriétés

Related Categories	Analytical/Chromatography, Astec CYCLOBOND Chiral HPLC Columns (Native and Derivatized Cyclodextrins), CYCLOBOND I 2000 DMP, Chiral HPLC Columns, HPLC Plus...
packaging	pkg of 1 ea
application(s)	HPLC: suitable
L × I.D.	10 cm × 4.6 mm
matrix	silica particle platform
matrix active group	cyclodextrin, beta- phase
particle size	5 µm
pore size	100 Å
suitability	suitable for L45 per USP
mode of chromatography	chiral

montrer moins de résultats

Description

General description
The reaction of the 3,5-dimethylphenyl isocyanate with the hydroxyl groups of β-cyclodextrin results in a pi-basic phase similar in character to the naphthylethyl carbamate phases. The selectivity is greater for the CYCLOBOND I 2000 DMP when the chiral center of the analyte is part of a ring structure or is on the α carbon. This phase has been very useful for derivatized amines, like amphetamine ACQ.

• Bonded phase: 3,5-Dimethylphenyl carbamate modified β-cyclodextrin

Recommended products
Discover LiChropur reagents ideal for **HPLC** or **LC-MS** analysis

Legal Information
Astec is a registered trademark of Sigma-Aldrich Co. LLC
CYCLOBOND is a registered trademark of Sigma-Aldrich Co. LLC

Informations Sécurité

Aucune information relative à la sécurité de ce produit est actuellement disponible.

Documents

Certificat d'Analyse

Devis/ Commande de produits Bulk

FDS

Customers Also Viewed

 21107AST Astec® CYCLOBOND® I 2000 DMP Chiral HPLC Guard Cartridge, 5 µm particle size, L × I.D. 2 cm × 4 mm	 20662-U Thermogreen® LB-2 Septa, solid discs diam. 14 mm (9/16 in.), pkg of 50 ea	 20660-U Thermogreen® LB-2 Septa, solid discs diam. 12.5 mm (1/2 in.), pkg of 50 ea	 20657-U Thermogreen® LB-1 Septa, solid discs diam. 10 mm (13/32 in.)	 20653-U Thermogreen® LB-2 Septa, solid discs diam. 10 mm (13/32 in.), pkg of 50 ea
--	---	--	---	--

Recently Viewed

 20662-U Thermogreen® LB-2 Septa, solid discs diam. 14 mm (9/16 in.), pkg of 50 ea	 20660-U Thermogreen® LB-2 Septa, solid discs diam. 12.5 mm (1/2 in.), pkg of 50 ea	 20657-U Thermogreen® LB-1 Septa, solid discs diam. 10 mm (13/32 in.)	 20653-U Thermogreen® LB-2 Septa, solid discs diam. 10 mm (13/32 in.), pkg of 50 ea	 20644-U Supeltext® M-2 Packed Column Ferrule, 1/16 in. Column O.D. I.D. 1/16 in., 1/16 in. O.D. Column, pkg of 10 ea
--	---	---	--	--

Protocoles et articles

Related Content

Supelco HPLC Calculator for Isocratic or Gradient Method Transfer

Calculates conditions for transfer of an isocratic or gradient method from one HPLC column to another. Allows method scaling from microbore through preparative column range. Calculates time and solve...
Keywords: High performance liquid chromatography

Documentation référencée

Did you use this product in your Paper? If so click here.

read abstract

Separation of chiral furan derivatives by liquid chromatography using cyclodextrin-based chiral stationary phases.

Xinxin Han et. al

Journal of chromatography. A, 1063(1-2), undefined (2005-2-11)

The enantiomeric separation of a set of 30 new chiral furan derivatives has been achieved on native and derivatized beta-cyclodextrin stationary phases using high performance liquid chromatography

(HPLC). The hydroxypropyl-beta-cyclodextrin (Cyclobon...[Read More](#)

Reprint of: Enantiomeric separation of functionalized ethano-bridged Troger bases using macrocyclic cyclofructan and cyclodextrin chiral selectors in high-performance liquid chromatography and capillary

electrophoresis with application of principal component analysis Weatherly, Choyce A., et al. Journal of Chromatography. B, Biomedical Applications 968, 40-48, (2014)

Enantiomeric separation of functionalized ethano-bridged Troger bases using macrocyclic cyclofructan and cyclodextrin chiral selectors in high-performance liquid chromatography and capillary

electrophoresis with application of principal component analysis Weatherly, Choyce A., et al. Journal of Chromatography. B, Biomedical Applications 955-956, 72-80, (2014)

read abstract

Enantiomeric separations of illicit drugs and controlled substances using cyclofructan-based (LARIHC) and cyclobond I 2000 RSP HPLC chiral stationary phases.

Nilusha L T Padivitage et. al

Drug testing and analysis, 6(6), undefined (2013-10-12)

Recently a novel class of chiral stationary phases (CSPs) based on cyclofructan (CF) has been developed. Cyclofructans are cyclic oligosaccharides that possess a crown ether core and pendent fructofuranose moieties. Herein, we evaluate the applicabil...[Read More](#)

read abstract

The role of pi-acidic and pi-basic chiral stationary phases in the high-performance liquid chromatographic enantioseparation of unusual beta-amino acids.

István Ilisz et. al


Chirality, 21(3), undefined (2008-6-17)

The application of 3,5-dimethylphenyl-carbamoylated-beta-cyclodextrin (Cyclobond I 2000 DMP) and 2,6-dinitro-4-trifluoromethylphenyl-ether-beta-cyclodextrin-based (Cyclobond DNP) chiral stationary phases for the high-performance liquid chromatographi...[Read More](#)

Produits associés



Brighter LC-MS
LC-MS Grade Solvents and Reagents for accurate, brilliant results
[LEARN MORE](#)



Milli-Q® water purification solutions
Ultrapure & Pure Systems for Every Lab
[Learn More](#)