

TspDTI

5'-ATGAA(N)₁₁-3'
3'-TACTT(N)₉-5'

Cat. No.	size
E2502-01	50 units
E2502-02	250 units

Reaction Temperature: 70°C

Inactivation Temperature (20 min): —

Prototype: TspDTI

Source: *Thermus species DT*

Purified from *E.coli* strain that carries the cloned tspDTRI gene from *Thermus sp. DT*.

Package Contents:

- TspDTI
- 10 x Reaction Buffer TspDTI

Storage Conditions: Store at -20°C.

Prepare and store buffer aliquots at -70°C.

DNA Methylation:

No Inhibition: dam, dcm, EcoKI, CpG

Standard Reaction Protocol (for 30 µl volume):

Mix the following reaction components:

1-2 µg pure DNA or 10 µl PCR product (≈0.1-2 µg DNA)

3 µl 10 x Buffer TspDTI

1-2 U TspDTI (use 1 U per µg DNA, < 10% React. Volume!)

Tips: Add enzyme as last component. Mix components well before adding enzyme. After enzyme addition, mix gently by pipetting. Do not vortex.

add sterile H₂O to 30 µl final volume

Incubate for 3 h at 70°C

Stop reaction by alternatively

(a) Addition of 1.2 µl EDTA pH 8.0 [0.5 M], final 20 mM or

(b) Heat Inactivation

(not applicable for this enzyme) or

(c) Spin Column DNA Purification

(e.g. EURx PCR/DNA Clean-Up Kit, Cat.No. E3520) or

(d) Gel Electrophoresis and Single Band Excision

(e.g. EURx Agarose-Out DNA Kit, Cat.No. E3540) or

(e) Phenol-Chloroform Extraction or Ethanol Precipitation.

Note 1: It is required to purify DNA before digestion. We recommend PCR/DNA Clean-Up Purification Kit or Agarose-Out DNA Purification Kit.

Note 2: It is not recommended to use more than 2 units per 30 µl reaction. It is strongly suggested to perform digestion for over 1 hr.

Note 3: To avoid DNA shift during electrophoresis caused by strong protein-DNA interaction, it is recommended to terminate reaction by addition of reaction stop solution (containing denaturing reagent, i.e. 0.2% SDS) followed by 20 minutes heat inactivation in 89°C.

Unit Definition:

One unit is the amount of enzyme required to digest 1 µg of pUC19 DNA to obtain stable digestion pattern in 1 hr in a total reaction volume of 30 µl. Enzyme activity was determined in the recommended reaction buffer.

Reaction Buffer:

1 x TspDTI Buffer: 10 mM Tris-HCl (pH 8.5 at 25°C), 10 mM MgCl₂, 1 mM dithiothreitol + enhancers.

Avoid multiple cycles of freezing/thawing of the stock reaction buffer (no more than 3 times). Thawing should be performed at temperatures not exceeding 10°C. Recommended procedure is to divide the provided reaction buffer into smaller portions and preserve them at -70°C for long-term. Temperature of -20°C should be used only for short-term storage.

Storage Buffer:

20 mM Tris-HCl (pH 8.3 at 25°C), 25 mM (NH₄)₂SO₄, 25 mM KCl, 0.5 mM EDTA, 0.5 mM dithiothreitol, 0.02% Triton™X-100, 0.02% Tween™20, 0.02% Igepal, 50% (v/v) glycerol.

Quality Control:

All preparations are assayed for contaminating endonuclease, 3'-exonuclease, 5'-exonuclease/5'-phosphatase, as well as for nonspecific single- and double-stranded DNase activities.