

Toutes les
photos (1)

Documents

 [FDS](#)

 [COO/COA](#)

11034731001 Roche

Primer "random"

lyophilized, sufficient for ≤ 400 reactions, pkg of 2 mg (Primer, Random pd(N)6 Potassium Salt)

Synonyme(s):


primer

Référence	Conditionnement	Disponibilité	Prix	Quantité
11034731001	2 MG	 Only 5 left in stock (more on the way)	Détails... 329,00 €	- + 

[Demander une commande en gros](#)

[Ajouter au panier](#)

PRODUITS RECOMMANDÉS



Sigma-Aldrich

H0268

Hexanucleotide Primers

lyophilized powder



Roche

10814270001

Primer for cDNA Synthesis

lyophilized, pkg of 40 μ g

Consulter le prix et la disponibilité

Consulter le prix et la disponibilité

PROPRIÉTÉS

Forme	lyophilized
Niveau de qualité	100
Utilisation	sufficient for ≤400 reactions
Conditionnement	pkg of 2 mg (Primer, Random pd(N)6 Potassium Salt)
manufacturer/tradename	Roche
Expédié(e)s dans	dry ice
Temp. de stockage	−20°C

Catégories apparentées

Nucleic Acid Labeling

Roche® Life Science Products

DESCRIPTION

Description générale

p(dN)₆

Application

Primer "random" has been used:

- in a reverse transcription-polymerase chain reaction (RT-PCR) and quantitative PCR (qPCR) to retro-transcribe RNA
- to reverse-transcribe RNA samples to cDNA
- in digital droplet PCR (ddPCR) to synthesis cDNA from RNA

Primer for cDNA synthesis

Caractéristiques et avantages

- Offers a consistent representation of all RNA sequences in mRNA
- Can prime cDNA transcription from RNAs lacking a poly(A) tail
- Short cDNA transcripts can overcome the challenges posed by RNA secondary structures
- The average length of cDNAs formed can be controlled by adjusting the ratio of random primers to RNA in the reverse transcription (RT) reaction

Autres remarques

For life science research only. Not for use in diagnostic procedures.

INFORMATIONS SUR LA SÉCURITÉ

Code de la classe de stockage

12 - Non Combustible Liquids

WGK

nwg

Flash Point(F)

does not flash

Point d'éclair C

does not flash

DOCUMENTATION

Certificat d'analyse

Saisir un numéro de lot pour rechercher un certificat d'analyse (COA).

Numéro de lot

e.g. 023J5431

Comment saisir un numéro de lot (COA)

Rechercher

Certificat d'origine

Saisir un numéro de lot pour rechercher un certificat d'origine (COO).

Numéro de lot

e.g. 023J5431

Comment saisir un numéro de lot (COO)

Rechercher

LES CLIENTS ONT ÉGALEMENT CONSULTÉ



Roche

TRANSRTRO

Transcriptor Reverse Transcriptase

[Consulter le prix et la disponibilité](#)



Roche

11483188001

First Strand cDNA Synthesis Kit for RT-PCR (AMV)

sufficient for 30 reactions (including 5 control reactions), kit of 1 (10 components), suitable for RT-PCR, hotstart...

[Consulter le prix et la disponibilité](#)

ARTICLES REVUS PAR DES PAIRS

Novel RNA-binding activity of MYF5 enhances Ccnd1/Cyclin D1 mRNA translation during myogenesis.

Amaresh C Panda et al.

Nucleic acids research, 44(5), 2393-2408 (2016-01-29)

Skeletal muscle contains long multinucleated and contractile structures known as muscle fibers, which arise from the fusion of myoblasts into multinucleated myotubes during myogenesis. The myogenic regulatory factor (MRF) MYF5 is the earliest to be expressed during myogenesis and functions

Large-scale phenome analysis defines a behavioral signature for Huntington's disease genotype in mice.

Vadim Alexandrov et al.

Nature biotechnology, 34(8), 838-844 (2016-07-05)

Rapid technological advances for the frequent monitoring of health parameters have raised the intriguing possibility that an individual's genotype could be predicted from phenotypic data alone. Here we used a machine learning approach to analyze the phenotypic effects of polymorphic

Histone demethylase JMJD2A drives prostate tumorigenesis through transcription factor ETV1.

Tae-Dong Kim et al.

The Journal of clinical investigation, 126(2), 706-720 (2016-01-06)

Histone demethylase upregulation has been observed in human cancers, yet it is unknown whether this is a bystander event or a driver of tumorigenesis. We found that overexpression of lysine-specific demethylase 4A (KDM4A, also known as JMJD2A) was positively correlated


Shedding a new light on Huntington's disease: how blood can both propagate and ameliorate disease pathology.

Marie Rieux et al.

Molecular psychiatry, 26(9), 5441-5463 (2020-06-10)

Huntington's disease (HD) is a monogenic neurodegenerative disorder resulting from a mutation in the huntingtin gene. This leads to the expression of the mutant huntingtin protein (mHTT) which provokes pathological changes in both the central nervous system (CNS) and periphery.

PRODUITS CONSULTÉS RÉCEMMENT



Roche

LEU-RO

Leupeptin

powder, from synthetic, =96.5%

[Consulter le prix et la disponibilité](#)



Roche

RPOLT3-RO

T3 RNA Polymerase

from *Escherichia coli* HB101

[Consulter le prix et la disponibilité](#)

SERVICE TECHNIQUE

Notre équipe de scientifiques dispose d'une expérience dans tous les secteurs de la recherche, notamment en sciences de la vie, science des matériaux, synthèse chimique, chromatographie, analyse et dans de nombreux autres domaines..

[Contacter notre Service technique](#)

Le contenu de cette page vous a-t-il été utile ?*



Que pouvons-nous faire pour améliorer cette page spécifique de notre site Internet ?

Envoyer

© 2022 Merck KGaA, Darmstadt, Allemagne et/ou ses sociétés affiliées. Tous droits réservés.

La reproduction d'une quelconque partie du contenu de ce site est strictement interdite sans autorisation.

[Conditions d'utilisation du site](#) | [Politique de confidentialité](#) | [Conditions générales de vente](#) | [Consentement relatif au copyright](#) | [Paramètres des cookies](#)