

#61977 Store at -20C

Human FGF-basic/FGF2 (154 aa) Recombinant Protein

10 µg



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

MW (kDa):
17

UniProt ID:
#P09038

Entrez-Gene Id:
2247

Background

FGF basic (FGF2) is produced by epithelial, tumor and other cell types (1). FGF basic is involved in developmental processes and regulates differentiation, proliferation, and migration (1-6). FGF basic is a critical factor for growing embryonic stem cells in culture without inducing differentiation. FGF basic has a high affinity for heparan sulfate (1,2) and FGF-heparan sulfate binding is a step in the activation of FGFR tyrosine kinase. There are four distinct FGF receptors and each has multiple splice variants. FGF basic binds with high affinity to many, but not all, FGFRs. Signaling cascades activated through FGF basic binding to FGFR include the ras-raf-MAPK, PLCγ/PKC, and PI3K/AKT pathways (1).

Endotoxin

Less than or equal to 1 EU / 1 µg hFGF-basic

Purity

A greater than 95% purity was determined by SDS-PAGE.

Source / Purification

Human FGF-basic Recombinant Protein was expressed in *E. coli* and is supplied in a lyophilized form.

Bioactivity

The bioactivity of Human FGF-basic Recombinant Protein was determined in a NIH/3T3 cell proliferation assay. The ED₅₀ of each lot is between 0.05 - 1.0 ng/ml.

Background

FGF basic (FGF2) is produced by epithelial, tumor and other cell types (1). FGF basic is involved in developmental processes and regulates differentiation, proliferation, and migration (1-6). FGF basic is a critical factor for growing embryonic stem cells in culture without inducing differentiation. FGF basic has a high affinity for heparan sulfate (1,2) and FGF-heparan sulfate binding is a step in the activation of FGFR tyrosine kinase. There are four distinct FGF receptors and each has multiple splice variants. FGF basic binds with high affinity to many, but not all, FGFRs. Signaling cascades activated through FGF basic binding to FGFR include the ras-raf-MAPK, PLCγ/PKC, and PI3K/AKT pathways (1).

Background References

1. Dvorak, P. and Hampl, A. (2005) *Folia Histochem Cytobiol* 43, 203-8.
2. Ornitz, D.M. and Itoh, N. (2001) *Genome Biol* 2, REVIEWS3005.
3. Shi, Y. et al. (2008) *Crit Rev Oncol Hematol* 65, 43-53.
4. Fontijn, D. et al. (2006) *Br J Cancer* 94, 1627-36.
5. Marek, L. et al. (2009) *Mol Pharmacol* 75, 196-207.
6. Acevedo, V.D. et al. (2009) *Cell Cycle* 8, 580-8.

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or

therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.