

# Active Recombinant Human FGF-2/bFGF Protein

Catalog No.: RP01042 **Recombinant**

## Sequence Information

Species	Gene ID	Swiss Prot
Human	2247	P09038-4

### Tags

No tag

### Synonyms

BFGF; FGF-2; FGFB;  
HBGF-2; FGF2; FGF-2; FGFB; HBGF-2; Basic FGF; BFGF; fibroblast growth factor 2

## Product Information

Source	Purification
<i>E. coli</i>	> 95% by SDS-PAGE.

### Endotoxin

< 1.0 EU/μg of the protein by LAL method.

### Formulation

Lyophilized from a 0.22 μm filtered solution of 20mM Tris, 150 mM NaCl, pH7.5. Contact us for customized product form or formulation.

### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## Background

### Basic Information

#### Description

Active Recombinant Human FGF-2/bFGF Protein is produced by *E. coli* expression system. The target protein is expressed with sequence (Pro143-Ser288) of human FGF2 (Accession #NP\_001997.5).

#### Bio-Activity


1. Measured by its binding ability in a functional ELISA. Immobilized Human FGF2 at 0.5 μg/mL (100 μL/well) can bind Human GPC3 with a linear range of 7-20 ng/mL. 2. Measured in a cell proliferation assay using BALB/c 3T3 mouse embryonic fibroblasts. The ED<sub>50</sub> for this effect is typically 0.635-2.54 ng/mL, corresponding to a specific activity of  $3.94 \times 10^5 \sim 1.57 \times 10^6$  units/mg. 3. Recombinant Human VEGFA (40 ng/mL, Cat. RP01162) and bFGF (50 ng/mL) induce mesoderm cells to differentiate into hematopoietic stem and progenitor cells. After 4 days induction, pebbly-like CD43+ hematopoietic stem and progenitor cells appeared in the hematogenic endothelium. 4. The primary neural stem cells were cultured with 20 ng/mL bFGF and observed every 24 h. Results showed that the particle size of the suspended neural stem cells gradually increased.

#### Storage


Store the lyophilized protein at -20°C to -80 °C for long term. After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week. Avoid repeated freeze/thaw cycles.

## Contact


## Validation Data




Recombinant Human FGF-2/bFGF Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 17 kDa.




Immobilized recombinant human FGF2 at 1 µg/mL (100 µL/well) can bind recombinant human FGFR4 with a linear range of 30-125 ng/mL.




Immobilized Human FGF2 at 0.5 µg/mL (100 µL/well) can bind Human GPC3 with a linear range of 7-20ng/mL.



Recombinant Human FGF-2 promotes the proliferation of BALB/c 3T3 mouse embryonic fibroblasts cells. The ED<sub>50</sub> for this effect is typically 0.635-2.54 ng/mL, corresponding to a specific activity of 3.94 × 10<sup>5</sup>~1.57 × 10<sup>6</sup> units/mg.



Recombinant Human VEGFA(40 ng/mL, Cat. RP01162) and bFGF(50 ng/mL) induce mesoderm cells to differentiate into hematopoietic stem and progenitor cells. After 4 days induction, pebbly-like CD43+ hematopoietic stem and progenitor cells appeared in the hematogenic endothelium.(Customer feedback data)



Primary neural stem cells were cultured with a final concentration of 20 ng/mL FGF2, and as shown in the figure, the size of the suspended neural stem cells gradually increased.(Customer feedback data)