

Active Recombinant Human PCSK9 Protein

Catalog No.: RP01148 Recombinant

Sequence Information

SpeciesGene IDSwiss ProtHuman255738Q8NBP7

Tags

C-His

Synonyms

FH3; HCHOLA3; LDLCQ1; NARC-1; NARC1; PC9;PCSK9; FH3; proprotein convertase subtilisin/kexin type 9;HCHOLA3;LDLCQ1;NARC-1;NARC1;PC

Product Information

Source	Purification
HEK293 cells	> 92% by SDS
	PAGE.

Endotoxin

< 0.1 EU/ μ g of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

•

www.abclonal.com

Background

Basic Information

Description

Active Recombinant Human PCSK9 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gln31-Gln692) of human PCSK9 (Accession #NP_777596.2) fused with a 6×His tag at the C-terminus.

Bio-Activity


Measured by its binding ability in a functional ELISA. Immobilized Human LDLR at $5\mu g/mL$ (100 $\mu L/well$) can bind Human PCSK9 with a linear range of 0.5-294.7ng/mL.

Storage


Store the lyophilized protein at -20°C to -80 °C for long term. After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Active Active Recombinant Human PCSK9 Protein was determined by SDS-PAGE with Coomassie Blue, showing bands at 66 kDa and 19 kDa.



Immobilized Human LDLR at $5\mu g/mL$ (100 $\mu L/well$) can bind Human PCSK9 with a linear range of 0.5-294.7ng/mL.