

# Recombinant Rat Lipocalin-2/NGAL/LCN2 Protein

Catalog No.: RP01225 Recombinant

### **Sequence Information**

**Species Gene ID Swiss Prot**Rat 170496 P30152

### Tags

C-His

### **Synonyms**

NGAL;LCN2;Lipocalin-2;24p3;MSFI;LCN

### **Product Information**

Source	Purification
HEK293 cells	> 95% by SDS
	DAGE

### **Endotoxin**

 $< 0.1 EU/\mu g$  of the protein by LAL method.

### **Formulation**

Lyophilized from a 0.22 µm filtered solution of 20mM Tris,500mM NaCl, pH 7.0Contact us for customized product form or formulation.

### Reconstitution

Centrifuge the vial before opening.
Reconstitute to a concentration of
0.1-0.5 mg/mL in sterile distilled water.
Avoid votex or vigorously pipetting the
protein. For long term storage, it is
recommended to add a carrier protein
or stablizer (e.g. 0.1% BSA, 5% HSA,
10% FBS or 5% Trehalose), and aliquot
the reconstituted protein solution to
minimize free-thaw cycles.

### Contact



www.abclonal.com

### **Background**

### **Basic Information**

#### Description

Recombinant Rat Lipocalin-2/NGAL/LCN2 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gln21-Asn198) of rat Lipocalin-2/NGAL (Accession #NP\_570097.1.) fused with a 6×His tag at the Cterminus.

#### **Bio-Activity**

Measured by its ability to bind Iron(III) dihydroxybenzoic acid [Fe(DHBA)3]. The binding of Fe(DHBA)3 results in the quenching of Trp fluorescence in Lipocalin2 . It binds >51330  $\mu$ M of Fe(DHBA)3.


#### Storage

Store the lyophilized protein at -20°C to -80 °C for long term.


After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week.

Avoid repeated freeze/thaw cycles.

## **Validation Data**



Recombinant Rat Lipocalin-2/NGAL/LCN2 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 27 kDa.



The binding of Fe(DHBA)3 results in the quenching of Trp fluorescence in Lipocalin2 . It binds >51330  $\mu$ M of Fe(DHBA)3.